4CPS-139 THE PYRAMID STUDY: PATIENTS TREATED BY MYELOMA ORAL MEDICATION HAVE A BAD COMPLIANCE

N Allaire*, JF Huon, P Nizet, C Dubreuil, V Burgeot, I Rouiller, L Flet, M Tching-Sin. *CHU Nantes, Pharmacy, Nantes, France*

10.1136/ejhp-2019-eahpconf.288

Background Like other cancerous diseases, there are now many oral treatments for haematological malignancies. These oral products have a lot of advantages, but their effectiveness depends on patient compliance. Do they have a good effect?

Purpose This study aimed to assess patient medication compliance for myeloma oral treatments (Imids: Thalidomid, Lenalidomid, Pomalidomid) and to identify factors associated with medication compliance by adult outpatients.

Material and methods A cross-sectional prospective study was conducted at a university teaching hospital from January to August 2018.

Investigational medication compliance was assessed at the time of the renewal dispensation using the Morisky questionnaire. Other questionnaires were completed by outpatients at home: SF-36 quality of life, SATMED-Q satisfaction of treatment and HLQ health literacy. Demographic and other baseline data were collected using self-administered questionnaires.

Results Of the 62 participants, 36 (58.1%) were male, and the median age was 69 years (range: 43–85 years). The mean compliance Morisky score was 7.12/8% and 45.9% of the patients had a score of 8/8 (good observance). Patients were divided into two groups for the analysis: the good adherence one (score >7, n=28) and the medium/poor adherence group (score ≤7, n=33).

In univariate analysis, the Morisky medication compliance score was negatively correlated with existence of a tip to remember drug intake ($p=0.0234$), being a member of a patients' association ($p=0.0118$) and positively correlated with treatment satisfaction ($p=0.0081$), perceived health ($p=0.0175$), a good psychic health ($p=0.0119$) and few limitations associated with it ($p=0.0047$).

Conclusion Only half the included outpatients had an optimal adherence of their medication. Having a poor satisfaction of their treatment and a poor psychic health were important risk factors for inadequate medication compliance.

This and other data suggest that hospitals and health professionals should encourage initiatives aimed at improving medication compliance in adults treated by Imids. Therapeutic education is not the easiest solution, due to the necessity of training. Pharmaceutical interviews could be easier to set up. To maximise the efficiency of these interventions, they should be aimed to poor compliance assumed patients.

REFERENCES AND/OR ACKNOWLEDGEMENTS

No conflict of interest.

4CPS-140 HOW PHARMACISTS CAN IMPROVE PATIENTS' CARE MANAGEMENT IN SUBCUTANEOUS ANTIPSORIATIC BIOLOGIC THERAPY

¹H Benoist*, ²A Dinahet, ²C Clement, ¹G Saint-Lorant. ¹Caen University Hospital, Pharmacy, Caen, France; ²Caen University Hospital, Dermatology, Caen, France

10.1136/ejhp-2019-eahpconf.289

Background Subcutaneous antipsoriatic biologic therapy (SCABT) facilitates ambulatory care. However, it requires a therapeutic education of patients (TEP) to enable patients to acquire the necessary skills for the proper use of medications and to ensure safe administration.

Purpose The aim of this study was to improve patients' care management on SCABT.

Material and methods This work took place in the dermatology unit (10 beds) of a university hospital. A patient and nurse survey was conducted during TEP sessions in order to identify requirements. TEP sessions took place directly after the dermatological consultation and were dedicated to one patient. A questionnaire evaluating nurses' knowledge on SCABT was distributed to each nurse (n=10). Following the implementation of pharmacist-led actions, a second round of the questionnaire was distributed. A feedback session of the nurses and a patient satisfaction survey were also done.

Results The survey of six patients showed that 100% did not read or still lose information supports from pharmaceutical companies. The nurses survey (n=6) had highlighted their need for adapted tools. The first round of the nurses (n=10) questionnaire showed less than 10% of correct answers regarding each SCABT characteristic. A nurse training course of 1 hour made by pharmacists was presented to nurses (n=5). This presentation contained all the essential information on SCABT. Leaflets of all SCABT were carried out in a multi-disciplinary approach to remind patients of all key points of SCABT. A dosing regimen on the leaflets allowed the patient to trace their home injections which were checked by the nurse on the following appointment. A poster 'Guide SCABT' was elaborated in order to sum up all information. The results of the nurse post-training questionnaire (n=4) was as follows: 100% of nurses knew methods of preservation, 100% dispensing modalities, 100% waste management and 50% SCABT characteristics. The 2 month feedback showed a general satisfaction of all patients (n=6) and all caregivers (n=7).

Conclusion This multidisciplinary approach helps meet patients' expectations and creates a dynamic and thorough TEP approach. It confirms that clinical pharmacy services help answer patients' and caregivers' needs.

REFERENCES AND/OR ACKNOWLEDGEMENTS

No conflict of interest.

4CPS-141 ECONOMIC IMPACT OVER THE PAST 4 YEARS ASSOCIATED WITH BIOLOGICAL THERAPY OPTIMISATION IN RHEUMATIC DISEASES PATIENTS

¹I Cavada Carranza*, ¹FJ Alonso-Zazo, ¹ME Martínez-Núñez, ¹R Vázquez-Sánchez, ¹J Sánchez-Rubio-Ferrández, ²Á Aragón-Díez, ²JM Rodríguez-Heredia, ¹T Molina-García. ¹Hospital Universitario de Getafe, Pharmacy Department, Madrid, Spain; ²Hospital Universitario de Getafe, Rheumatology Department, Madrid, Spain

10.1136/ejhp-2019-eahpconf.290

Background Therapeutic decision-making for biologic-therapies (BT) dose optimisation should be based on optimal disease activity results. The goal of BT optimisation is to guarantee long-term effectiveness and safety, maximising economic savings.

Purpose To evaluate BT optimisation patterns in patients with rheumatic diseases (RD) and associated economic savings.

Material and methods Observational and retrospective study.