4CPS-017

EVALUATION OF PHARMACISTS' INPUT IN ANTICOAGULATION CLINIC REVIEWING DIRECT ORAL ANTICOAGULANTS INITIATED IN A SECONDARY CARE HOSPITAL IN LONDON

A Grover*, P Wright, S Antoniou, C Jaggot. *Barts Health NHS Trust, Cardiology- Pharmacy, London, UK*

10.1136/ejhpharm-2019-eahpconf.166

Background The National Institute for Health and Care Excellence (NICE) technology appraisals has made recommendations on four direct oral anticoagulants (DOACs). Local anticoagulation policy recommends all patients newly initiated on a DOAC should be followed-up in an anticoagulation clinic within 4 weeks. There is evidence suggesting that up to 30% of patients are dosed inappropriately according to their age, bodyweight and renal function.

Purpose To assess the dosing appropriateness of DOACs at initiation at the 1 month follow-up anticoagulation appointment in the clinic.

- Determine percentage of patients who are initiated on an appropriate dose.
- Determine percentage of patients that had an intervention made in their treatment plan at the clinic.

Material and methods Data was collected retrospectively over a period of 6 months from patient healthcare records from January 2018 to July 2018 for patients attending anticoagulant clinics.

Results

- Sixty-four per cent (n=118) of patients newly initiated on a DOAC were followed-up within 4 weeks.
- Ninety-two per cent (n=166) of patients were initiated on an appropriate dose of DOAC in accordance with product licence.
- Fifteen per cent (n=27) of patients had either a dose or DOAC changed, or DOAC stopped at the follow-up appointment by a pharmacist.
- The majority of alterations were due to incorrect documentation of weight, use of old blood test results and use of eGFR instead of calculated creatinine clearance (CrCl) using Cockcroft and Gault.
- The majority of patients were followed up within a 4 week period. A significant proportion, 8% (n=17), required dose amendments, as initial dosing was incorrectly based on CrCL estimated by the hospital system which is based on e-GFR and not Cockcroft and Gault in line with the product licences and clinical trials.

Conclusion Pharmacists have a clear role in ensuring appropriate dosing of DOACs and a reminder (and education) for non-specialist pharmacists on the importance of dosing based on CrCl with Cockcroft and Gault, as opposed to the default on hospitals with is e-GFR.

REFERENCE AND/OR ACKNOWLEDGEMENTS

Manzoor BS, Cheng WH, Lee JC, *et al.* Quality of pharmacist-managed anticoagulation therapy in long-term ambulatory settings: a systematic review. *Ann Pharmacother* 2017;51:1122–37.

No conflict of interest.

4CPS-018

EVALUATION OF ANTIPLATELET AGENT PRESCRIBING IN PATIENTS ON DIRECT ORAL ANTICOAGULANT

¹AS Larock*, ^{2,3}A Spinewine, ⁴P Laloux, ⁵P Eucher, ⁶C Hanet. ¹CHU UCL Namur- Namur Thrombosis and Haemostasis Centre- Université Catholique de Louvain, Pharmacy, Yvoir, Belgium; ²CHU UCL Namur- Namur Thrombosis and Haemostasis Centre- Université Catholique de Louvain, Pharmacy, Yvoir, Belgium; ³Louvain Drug Research Institute- Clinical Pharmacy Research Group- Université Catholique de Louvain, Pharmacy, Brussels, Belgium; ⁴CHU UCL Namur- Université Catholique de Louvain, Neurology, Yvoir, Belgium; ⁵CHU UCL Namur- Université Catholique de Louvain, Cardiac- Vascular and Thoracic Surgery, Yvoir, Belgium; ⁶CHU UCL Namur- Université Catholique de Louvain, Cardiology, Yvoir, Belgium

10.1136/ejhpharm-2019-eahpconf.167

Background Among patients requiring an oral anticoagulant (OAC), a large proportion also take an antiplatelet agent (AP). Several studies have highlighted the significantly increased bleeding risk associated with a combined OAC (VKA mainly) and AP (aspirin mainly) therapy, without a reduction in risk of recurrence of coronary artery events or thromboembolism. The continuation of an AP in patients on OAC therapy for venous thromboembolism or atrial fibrillation remains a recurrent matter of debate and is still little studied in patients on direct OAC (DOAC).

Purpose Our main objective was to evaluate to what extent combined DOAC-AP therapy met recommendations of current guidelines. A secondary objective was to describe antithrombotic prescription schemes in patients on DOAC with a recent percutaneous coronary intervention (PCI).

Material and methods We performed an observational retrospective cohort study in a 450-bed teaching hospital. Among DOAC patients prospectively reviewed by a clinical pharmacist dedicated to anticoagulation between January and December