aeruginosa (ATCCVR 9027TM), *Candida albicans* (ATCCVR 10231TM), *Aspergillus brasiliensis* (ATCCVR 16404TM) and *Staphylococcus aureus* (ATCCVR 6538TM).

To perform the growth assay, trypticase soy agar (TSA) were used for *P. aeruginosa* and *S. aureus*, and sabouraud glucose agar (SAB) for *C. albicans* and *A. brasiliensis*.

The test was performed by taking a 1:1000 dilution of 1 g of topical resorcinol in a 0.1% Tween 80 and phosphate buffered saline solution and adding 100 µL of a suspension equivalent to 1×10^3 cfu/mL of every ATCC strain, which were inoculated in TSA or SAB. All tests were done in duplicate and medium lectures were made in 48 hours.

Results The ability of ATCC strains to growth in resorcinol formulation was confirmed under the study conditions. There was mean growth of 17×10^4 cfu/mL for *S. aureus* and 11×10^4 cfu/mL for *P. aeruginosa* in TSA. For *A. brasiliensis* and *C. albicans*, 1×10^4 cfu/mL and 2×10^4 cfu/mL were detected, respectively.

Conclusion and relevance The presented method shows a simplified way to test the microbiological viability of 15% topical resorcinol for quality control.

REFERENCES AND/OR ACKNOWLEDGEMENTS

No conflict of interest.

3PC-003 TERLIPRESSIN PH STABILITY FOR CONTINUOUS INFUSION

A Castro Balado*, J González-López, S Blanco-Dorado, EJ Bandín-Vilar, L García-Quintanilla, I Varela-Rey, M Busto Iglesias, M González-Barcia, I Zarra-Ferro. *Complejo Hospitalario Universitario De Santiago De Compostela, Hospital Pharmacist Service, Santiago De Compostela, Spain*

10.1136/ejhp-pharm-2020-eahpconf.50

Background and importance Terlipressin is a synthetic vasoconstrictor peptide similar to desmopressin, used to treat bleeding oesophageal varices and hepatorenal syndrome. Recent publications have shown greater efficacy via continuous infusion (CI) compared with intermittent bolus injection. This could be explained by its pharmacodynamic effect (<4 hours): in 24 hours, with injections every 6 hours, there may be ≥ 8 hours without pharmacological effect. Terlipressin is available in ampoules and is stable only at pH 3–4. It is not currently known if there is a variation in pH after dilution for continuous infusion, and its impact on stability.

Aim and objectives The objective of the study was to determine the pH variation after dilution of terlipressin in different diluents commonly used in clinical practice for administration as a CI.

Material and methods The diluents used were 0.9% NaCl (NS), 5% dextrose (D5W) and 3.3% dextrose–0.3% saline (DS). The initial pH measurement was performed with the commercial ampoule (8.5 mL) after reaching room temperature, as well as with the diluents separately. Subsequently it was diluted to 10, 20, 50, 100, 200, 250 and 500 mL. Triplicate pH measurements were made. The whole process was carried out at 23°C with a precision pH meter WTW inoLab pH level 1.

Results Initial pH values were: 3.94 ± 0.04 terlipressin, 5.62 ± 0.10 NS, 6.14 ± 0.04 D5W and 4.64 ± 0.04 DS. In NS, up to 10 and 20 mL, a slight decrease in pH was observed up to 3.92 ± 0.03 . Subsequently, the value increased exponentially, reaching a pH of 4.11 ± 0.06 in 500 mL of NS. This initial

behaviour was only observed with NS, while with D5W and DS the pH increase was exponential with the increase in volume, until reaching pH values of 4.15 ± 0.01 and 4.13 ± 0.02 , respectively, at 500 mL. In all three cases, a tendency to reach pH values >4 was observed, values in which the stability of the molecule would be compromised.

Conclusion and relevance The results show that pH values are within the terlipressin stability range. This makes it possible to dilute terlipressin with NS, D5W and DS in volumes between 50 and 500 mL (not higher), allowing administration in 24 hours by CI, reducing the dose and number of administrations.

REFERENCES AND/OR ACKNOWLEDGEMENTS

No conflict of interest.

3PC-004 PHYSICOCHEMICAL STABILITY OF CEFEPIME IN POLYPROPYLENE SYRINGES AND IN ELASTOMERIC DEVICES

¹E D'huart*, ¹H Zenier, ¹J Vigneron, ²A Charmillon, ¹B Demore. ¹University Hospital of Nancy, Pharmacy Department, Vandoeuvre Lès Nancy, France; ²University Hospital of Nancy, Infectious and Tropical Diseases Department, Vandoeuvre Lès Nancy, France

10.1136/ejhp-pharm-2020-eahpconf.51

Background and importance Cefepime is a fourth generation cephalosporin used to treat severe infectious. For β -lactam antibiotics, publications have demonstrated that continuous administration is the preferred mode of administration. To the best of our knowledge, no stability data for cefepime solutions at 110 mg/mL in polypropylene syringes or at 50 mg/mL in elastomeric devices have been published.

Aim and objectives The objectives of the study were to assess the stability of cefepime solutions (1) at 110 mg/mL, in 0.9% sodium chloride (0.9% NaCl) or 5% dextrose (D5W), in polypropylene syringes at 20–25°C and (2) at 50 mg/mL, in 0.9% NaCl, in elastomeric devices at 37°C, after preparation and after storage for 6, 24 and 48 hours.

Material and methods Three preparations for each condition were made. For each analysis, one sample was taken from each preparation and analysed by high performance liquid chromatography. The method was validated according to the International Conference on Harmonisation Q2 (R1). Physical stability was evaluated by visual and subvisual inspection with a nephelometer. pH values were measured.

Results

1. In syringes, for each solvent, cefepime solutions at 110 mg/mL retained more than 90% of the initial concentration after 24 hours. No visual modification and no turbidity were observed. After 48 hours, the solutions retained around 83% of the initial concentration and pH values increased with the addition of 1 pH unit compared with the initial value.
2. In elastomeric devices, cefepime solution in 0.9% NaCl at 50 mg/mL retained more than 90% of the initial concentration over a period of 6 hours. After 24 and 48 hours, the solutions retained around 83% and 59% of the initial value, respectively. After 6 hours, visual colour modifications were observed. Under this condition, the initial pH value was 4.81 and 6.18 after 24 hours.

Conclusion and relevance The stability of cefepime in 0.9% NaCl and D5W at 110 mg/mL was demonstrated for 24 hours in syringes at 20–25°C. These stability data provide additional