Are we offering patients the right medicines information? A retrospective evaluation of readability and quality in online patient drug information

Laia Robert Sabaté, Laura Diego

Catalan Drug Information Center, Barcelona, Spain

Correspondence to

Dr Laia Robert Sabaté, Catalan Drug Information Center, Barcelona 08017, Spain; lrobert@cedimcat.info

Received 23 September 2019

Revised 26 January 2020

Accepted 4 February 2020

Published Online First

17 February 2020

EAHP Statement 6: Education and Research.

ABSTRACT

Objectives The aim of this study was to assess the quality and readability of patient drug information published on a corporate website and written by a multidisciplinary team including pharmacists, doctors, nurses, journalists, experts in healthcare communication, and patients. Documents were user-tested with patients in order to improve and adapt the final product.

Methods Readability was measured using the INFLESZ tool, a software developed from the Flesch formula and adapted to Spanish. DISCERN and EQIP were used for quality appraisal by two independent raters as they have shown to be useful and consistent for assessing information related to treatment and treatment choices.

Results Most of the documents (>67%) had an easy readability level, implying that they would be easily understood by a person with a primary school education level. The DISCERN tool showed higher reliability and concordance than EQIP. The overall DISCERN mean score for the documents was 55.4% for rater 1 and 51% for rater 2, implying very good quality.

Conclusions This retrospective analysis supports the implemented workflow of the multidisciplinary team and the user-testing process and encourages continuation of this systematic development of documents. Although both EQIP and DISCERN are useful and widely used tools, according to our results we would favour DISCERN to evaluate patient drug information.

INTRODUCTION

The availability of online healthcare information allows access to an unlimited number of resources. A recent Spanish survey, however, shows that physicians and pharmacists remain the most frequent source of health information (88.7% and 67.1%, respectively) for patients. The internet as a source of online health information is growing and approximately 60% of internet searches relate to healthcare queries.¹ Searches cover a wide range of diseases, medical treatments, and lifestyle issues such as diet, nutrition, or exercise.

Over the past two decades, the role of the patient has shifted from a doctor-patient paternalistic relationship to a partnership with shared care, shared decision making, and shared responsibilities.² Healthcare systems are now working towards a model where clinicians and patients work together to reach a decision about an intervention based on clinical evidence and patient preference. This process can be applied to many types of healthcare decisions, including whether to take a medication

or not.^{3,4} Information regarding medications plays a key role in helping patients make informed choices.

The Catalan Department of Health has implemented a new healthcare social media policy in Catalonia which involves offering evidence-based, reliable, and practical information about drugs and pharmacotherapy to patients and citizens. A multidisciplinary team, involving not only pharmacists (primary care, hospital, and community pharmacists), doctors, and nurses but also journalists and other experts in healthcare communication, was built to implement this new strategy. Their main tasks are to identify patient information needs and topics of interest and produce written drug information aligned with local medicine optimisation policies. Each new document produced is reviewed and validated by patients: this improves the final product before being published as the information has been tested by the final user and required amendments have been made. A questionnaire and short semi-structured interview was used for this user-testing, in order to assess whether people can find and understand key points of information. Every document has been tested in two rounds consisting of six to eight people.

Medicines information, if well designed and written, can enhance people's knowledge and contribute to informed decision making about their medication.⁵ Thus, the team set the objective in their ongoing improvement plan to assess the quality and readability of drug information published on the corporate website (<http://medicaments.gencat.cat/>) since 2015.

METHODS

Readability and quality were considered the key elements to appraise. A search was conducted in order to select the most suitable instruments developed for this task. Various tools/instruments and computerised formulas are available to assess readability including the Fry formula, SMOG, and Flesch-Kincaid. These instruments are mainly developed in English but some have been adapted to Spanish.⁶ These formulas are based on the average length of words and sentences and do not take into account other factors that may facilitate reading. They do not provide information on reliability, structure, or layout, which is their main drawback. Thus, other tools are required to assess the quality and accessibility of drug information.⁷ There are a variety of validated instruments available to assess the quality of written patient drug information (SAM, PEMAT, BALD, IDAPS, MIDAS, CIRF, ELF,

© European Association of Hospital Pharmacists 2021. No commercial re-use. See rights and permissions. Published by BMJ.

To cite: Robert Sabaté L, Diego L. *Eur J Hosp Pharm* 2021;**28**:144–148.

DISCERN, EQUIP).^{8–12} Certain tools, such as MIDAS, CIRF, and ELF15,⁹ were specifically developed for rating medication leaflets while other instruments, such as JAMA benchmarks or Health on the Net, rate the quality of health website information.^{13–14} We decided to use DISCERN and EQUIP, as these tools have shown usefulness and consistency in the quality assessment of information related to treatment and treatment choices.^{11–12}

Sixty documents published in the past 5 years on the Health Department website were included in this study. The documents covered a variety of topics ranging from drug treatment options, safety issues, best practices, and general information about medicines.

INFLESZ, an online software developed by Barrio Catalejo in 2008, was used for the readability analysis.¹⁵ INFLESZ was chosen as it is the model tool developed from the Flesch formula and is adapted to the average Spanish reader. It is routinely used in healthcare to assess the readability of informed consent documents, medicine leaflets, and reading materials for health education purposes.^{15–17} INFLESZ classifies the readability of the materials into five groups: “very difficult”, “difficult”, “normal”, “easy,” and “very easy”.

For the quality analysis, two independent raters evaluated the material. Both raters had medical information backgrounds and were given a set of DISCERN and EQUIP instruments as well as guidelines for their use. The first step was to reach a consensus regarding the criteria to apply in order to overcome the certain ambiguities the documents may carry and that could interfere with the assessment process.

DISCERN can be used by both patients and health professionals and is a validated instrument designed to assess the appropriateness of written health information for treatment choices. Since its development in the UK in 1999, DISCERN has been used in several studies to assess patient information on cancer, chronic illnesses, or ADHD.^{18–20} This instrument comprises a set of 16 questions. The first eight questions address document reliability, the following seven questions address specific items related to treatment choices, and the last question is the overall rate.¹¹ It includes open questions such as the aim of the document, topic relevance, accuracy and updates, treatment options and their impact on quality of life, as well as the benefits and side effects of therapies. The DISCERN score is based on a five-point Likert scale ranging from “no” (1 point) to “yes” (5 points). DISCERN scores range from 0 to 80 where the highest score represents a high-quality or excellent document. The DISCERN scale classifies readability as follows: poor ($\leq 15\%$), fair (16%–31%), good (32%–47%), very good (48%–63%), and excellent (64%–80%).

EQUIP is another validated instrument used by health professionals and researchers to measure the quality of any written health material (both printed and online). It is a 20-item tool and unlike DISCERN, EQUIP also evaluates aspects of design and language.¹² This instrument, developed for use in the paediatric setting, takes into account relevant informational needs for parents and caretakers as well.¹² It has been used for rating patient information on topics such as bariatric surgery, eczema, diabetes, or statin prescribing in primary care.^{21–24} The EQUIP score is calculated by adding one point when answering yes, 0.5 where the answer is “partially”, and zero points when the answer is “no”. Overall punctuation is expressed by percentage, the higher the percentage, the better the quality. Based on the score, EQUIP authors make one of the following recommendations: >75% maintain publication, review in 2 to 3 years; 51%–75% maintain publication, review in 1 to 2 years; 26%–50% maintain publication, start editorial review immediately and replace within 6 months to 1 year; and 0%–25% unpublish immediately.

Table 1 Readability results using the INFLESZ index

INFLESZ	Punctuation	Section	Publication's type	Results	
				Frequency	%
0–40		Very difficult	University, scientific	5	8.2
40–50		Difficult	Bachelor's degree, scientific dissemination, specialised press	13	21.3
55–65		Normal	Secondary school, general press, sports press	0	0.0
65–80		Easy	Primary school, tabloids, best-seller books	41	67.2
>80		Very easy	Primary school, comics	2	3.3

STATISTICAL ANALYSIS

Data analysis was performed using SAS software v9.4, SAS Institute Inc., Cary, NC, USA. A P-value < 0.05 was considered statistically significant.

Agreement between the results of the two raters was measured using Cohen's Kappa and the intra-class correlation coefficient (ICC). Reliability was assessed using Cronbach's alpha coefficient. The correlation between scores for each instrument was calculated using Pearson's correlation.

RESULTS

Readability analysis

In this study the mean INFLESZ index was 71.96 (SD 4.7) which implies that the documents were easy to read. Any result above 55 points using this tool is considered easy to read. Although most documents (>67%) had an easy readability level, a few documents were rated as difficult or very difficult to read (21.3% and 8.2% respectively). Table 1 displays the results of the INFLESZ index according to the five groups and provides a comparison with the education level and reading grade of Spanish publications.

Concordance analysis (Kappa coefficients and intra-class correlation coefficient)

Concordance was defined as the degree of agreement between the two raters and was calculated for each scale. Table 2 shows the results of the agreement analysis (using the Kappa coefficient to adjust for the effect of chance on the concordance observed) for both instruments including all items. Although the grade of agreement was considered high for both DISCERN and EQUIP, the kappa for DISCERN was remarkably lower than that for EQUIP (fair and good agreement, respectively).

ICC was calculated for the overall score of each instrument. ICC, defined as the proportion of total variability that is due to the variability of the subjects, allows assessment of the general agreement between two or more measurement or observation methods based on a variance analysis model (ANOVA) with repeated measures. A high concordance was seen with both instruments: DISCERN-ICC was 0.81 (95% CI 0.70–0.88) and EQUIP-ICC was 0.78, (95% CI 0.66–0.86). Values above 0.75 are considered to have excellent concordance.

Reliability and correlation analysis between raters

Reliability was defined as the ability of an instrument to measure consistently and was calculated using the Cronbach alpha coefficient. The closer the value of Cronbach alpha is to 1, the greater the internal consistency of the instrument. As seen in table 3, DISCERN showed higher reliability (Cronbach alpha coefficient > 0.77) than EQUIP for both raters.

Table 2 Kappa coefficients and ICC for DISCERN and EQIP

DISCERN				EQIP			
Item		K	Agreement (%)	Item		k	Agreement (%)
1	Aims described	0.18	67.2	1	Description of medical topic	0.33	72.1
2	Aims achieved	0.5	85.2	2	Tecnicismes	0.25	91.8
3	Relevance	0.34	80.3	3	Length of sentences	0.89	95.1
4	Sources of information	0.7	80.3	4	Addresses to the readers	0.4	67.2
5	Date of publication	0.89	31.1	5	Tone	1	100.0
6	Balance/bias	0.29	88.5	6	Design	0.66	98.4
7	Support/other resources	0.71	83.6	7	Illustrations or diagrams	0.24	47.5
8	Uncertainties	0.03	19.7	8	Order	0.16	88.5
9	Treatment description	0.08	29.5	9	Space for notes	1	100.0
10	Benefits	−0.04	27.9	10	Contact information	1	100.0
11	Risks	0.67	70.5	11	Date of publication	1	100.0
12	Results of non-treatment	0.27	50.8	12	Author	0.65	96.7
13	Quality of life	0.2	68.9	13	Implication of patients	0.93	96.7
14	Alternatives	0.53	73.8	14	Generic names	0.29	70.5
15	Support for shared decision making	0.28	47.5	15	Quality of life	0.68	78.7
				16	Support/other sources	0.88	83.6
				17	Purpose	0.55	91.8
				18	Benefits	0.44	78.7
				19	Risks	0.56	75.4
				20	Alternatives	0.69	82.0
Average		0.38	76.2	Average		0.63	85.7
ICC		0.81	(95% CI 0.70–0.88)	ICC		0.78	(95% CI 0.659–0.861)

For the EQIP scale, the mean score was 44% for both raters and thus the quality needed to improve, and documents should be reviewed in 6 months to 1 year (table 3). In contrast to these results, the overall mean score for the documents in the DISCERN scale was 55.4% for rater 1 and 51% for rater 2, which is considered very good in both instances.

Correlation between both instruments was calculated using Pearson's coefficient (r). A value of $r=1$ indicates perfect positive correlation. The strength of association between DISCERN and EQIP for both raters was 0.85 and 0.69 for rater 1 and 2, respectively, meaning a moderate-strong correlation.

DISCUSSION

The readability of more than two-thirds of the documents developed were considered easy to understand by a person with primary school level education. The readability level of the documents is similar to that of tabloids or best-seller books. Nevertheless, about one-third of the documents were rated as difficult or very difficult to read. This could be considered an accessibility barrier for certain citizens and should be reviewed in order to improve the overall quality of the documents produced. These results are slightly better than those from other studies where issues with online Spanish healthcare information readability

have been reported.^{25 26} Nevertheless, readability tools are considered informative as they count words and the length of sentences, but experts recommend to use them with other techniques to evaluate quality and scientific evidence.⁸

We used the EQIP and DISCERN tools for quality analysis as they are validated and have been used to assess information on patient medicine and drug choices. These instruments were developed in English, and we have not identified any similar tool to assess quality that has been validated in Spanish or Catalan. Our results show higher reliability and concordance with DISCERN when used to evaluate the quality of online patient drug information. This has also been observed in other studies.^{22 27}

Although global agreement calculated using ICC is higher for DISCERN, EQIP shows better mean Kappa values for each individual item evaluated in the instrument. The fact that the DISCERN five-point scale “forces” the rater to score even if an item does not apply may explain this difference. In EQIP there is an answer (“not apply”) that provides the opportunity to circumscribe the answers to only the items directly involved. Another factor that may have contributed is different standards between raters in terms of layout and document design that can be interpreted differently depending on the way the information is presented (printable PDF format or standard website html format).

Another limitation seen in our study is that results are only applicable to certain types of documents from the institutional website written in Catalan and Spanish and cannot be extrapolated to all patient drug information accessible online.

Although the analysis indicates that EQIP is a useful tool and addresses items such as layout and language, we would favour the use of DISCERN based on the results. This tool is especially useful for assessing documents or information developed to support shared decision making concerning drugs.

This retrospective analysis supports the implemented workflow and encourages the systematic development of documents

Table 3 Cronbach alpha coefficient and mean punctuation of EQIP and DISCERN

		EQIP	DISCERN
Rater 1	Cronbach alpha coefficient	0.6922	0.7793
	Mean	44.28	55.36
	Scale range	20.31–52.94	38–69
Rater 2	Cronbach alpha coefficient	0.4581	0.7916
	Mean	44.83	50.9
	Scale range	22.72–51.47	27–68

implemented 5 years previously. Nonetheless, in light of the EQIP results, the multidisciplinary team has decided to review the material annually.

The strength of this paper is that the documents were written by a multidisciplinary team involving doctors, nurses, pharmacists, civil servants, journalists, and professionals trained in communication. Thus, they incorporated a wider and more realistic vision of the healthcare system. This approach allows the detection of information needs, selection of topics of most interest to patients, and production and validation of documents until they are fit for purpose. As patient involvement in the development of drug information is considered key,²⁸ we decided to use a user-testing method. Pioneered by Professor David Sless in Australia in the 1990s, user-testing is a specific method to assess whether people can find and understand key information which can markedly improve performance.²⁹

Information about medicine that is well written, unbiased, reliable, easy to understand, and practical can increase people's knowledge and comprehension. This will enable patients:^{5 30}

- ▶ To improve pharmacotherapy literacy (individual's ability to obtain, evaluate, calculate, comprehend, and properly act on patient-specific information concerning pharmacotherapy and pharmaceutical services necessary to make appropriate medication-related decisions, regardless of the mode of content delivery)
- ▶ To use drugs safely and effectively
- ▶ To make informed decisions about medicines and their treatments.

CONCLUSION

The institutional web aims to encourage patients' self-care, promote rational drug use, and responsible utilisation of healthcare services. It offers scientific, objective, and updated information about drugs and treatments for citizens. Readability and quality appraisal are key to the continuous improvement plan and in order to offer scientific, objective, and updated information about drugs and treatments.

The analysis confirmed good readability and good quality of documents produced over the past few years by a multidisciplinary team (comprised of doctors, nurses, pharmacists, public administration agents, journalists, and experts in health communication) that have also been validated by patients. Although both EQIP and DISCERN are useful and widely used tools, according to our results we would favour DISCERN to evaluate patient drug information.

What this paper adds

What is already known on this subject

- ▶ Readability and quality of patient drug information are key elements to appraise in order to know if the information is fit for purpose.
- ▶ EQIP and DISCERN are validated and useful tools in the quality assessment of patient information about treatment and treatment choices.

What this study adds

- ▶ A multidisciplinary team (pharmacists, doctors, nurses, journalists, and other experts in healthcare communication) and a user-testing process have to be considered to develop quality patient drug information.

For those organisations interested in developing patient information concerning medicines, drug choices, and treatments we would recommend the workflow of the multidisciplinary team and the user-testing process to ensure documents are "fit for purpose".

Funding The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

Competing interests None declared.

Patient consent for publication Not required.

Provenance and peer review Not commissioned; externally peer reviewed.

Data availability statement Data are available upon request.

REFERENCES

- 1 Los ciudadanos ante la e-Sanidad. Opiniones Y expectativas de Los ciudadanos sobre El uso Y La aplicación de las tic en El ámbito sanitario. Observatorio nacional de las telecomunicaciones Y de la Sociedad de la Información. Available: <https://www.ontsi.red.es/ontsi/es/estudios-informes/los-ciudadanos-ante-la-e-sanidad-opiniones-y-expectativas-de-los-ciudadanos-sobre-> [Accessed 17 Jan 2019].
- 2 Coulter A. Paternalism or partnership? *BMJ* 1999;319:719–20.
- 3 Coulter A, Collins A. Making shared decision-making a reality: no decision about me, without me [Internet]. King's Fund, 2011. Available: <https://www.kingsfund.org.uk/publications/making-shared-decision-making-reality> [Accessed 12 Jun 2019].
- 4 Drug and Therapeutics Bulletin. An introduction to patient decision AIDS. *BMJ* 2013;347:f4147.
- 5 Raynor DKT. Written information on medicines for patients: learning from the PIL. *Drug Ther Bull* 2018;56:145–9.
- 6 Tip 6. Use Caution With Readability Formulas for Quality Reports. Content last reviewed May 2015. Agency for Healthcare Research and Quality, Rockville, MD. [Internet]. Available: <https://www.ahrq.gov/talkingquality/resources/writing/tip6.html> [Accessed 18 Jun 2019].
- 7 Guide to Appraisal Health Information. Patient Information Forum. For everyone involved in health information and support [Internet]. Available: https://www.pifonline.org.uk/wp-content/uploads/2014/11/Guide_AppraisingHealthInformation.pdf
- 8 Luk A, Aslani P. Tools used to evaluate written medicine and health information. *Health Educ Behav* 2011;38:389–403.
- 9 Mira JJ, Lorenzo S, Pérez-Jover V, et al. Assessment of the quality of medication information for patients in Spain. *Expert Opin Drug Saf* 2013;12:9–18.
- 10 Doack CC, Doack L, Roor J. Teaching Patients with Low Literacy Skills. 2nd ed. Philadelphia: JB Lippincott Company; 1996 [Internet]. 1996. Available: <https://www.hsph.harvard.edu/healthliteracy/resources/teaching-patients-with-low-literacy-skills/> [Accessed 7 May 2019].
- 11 Charnock D, Shepperd S, Needham G, et al. DISCERN: an instrument for judging the quality of written consumer health information on treatment choices. *J Epidemiol Community Health* 1999;53:105–11.
- 12 Moul B, Franck LS, Brady H. Ensuring quality information for patients: development and preliminary validation of a new instrument to improve the quality of written health care information. *Health Expect* 2004;7:165–75.
- 13 Silberg WM, Lundberg GD, Musacchio RA. Assessing, controlling, and assuring the quality of medical information on the Internet: caveat lector et viewer – let the reader and viewer beware. *JAMA* 1997;277:1244–5.
- 14 HONcode Principles. The Health On the Net Foundation [Internet]. Available: <https://www.hon.ch/en/certification.html#principles> [Accessed 22 May 2019].
- 15 Barrio-Cantalejo IM, Simón-Lorda P, Melguizo M, et al. Validación de la Escala INFLESZ para evaluar La legibilidad de Los textos dirigidos a pacientes. *An Sist Sanit Navar* 2008;31:135–52.
- 16 Mariscal-Crespo MI, Coronado-Vázquez MV, Ramírez-Durán MV. Análisis global de la legibilidad de los documentos de consentimiento informado utilizados en los hospitales públicos de España. *Revista de Calidad Asistencial* 2017;32:200–8.
- 17 Ballesteros-Peña S, Fernández-Aedo I. Legibility analysis of drug package leaflets using Flesch–Szigriszt index and INFLESZ score. *An Sist Sanit Navar* 2013;36:397–406.
- 18 Kaicker J, Borg Debono V, Dang W, et al. Assessment of the quality and variability of health information on chronic pain websites using the DISCERN instrument. *BMC Med* 2010;8:59.
- 19 Montoya A, Llopis N, Gilaberte I. Validation of the translation of an instrument to measure reliability of written information on treatment choices: a study on attention deficit/hyperactivity disorder (ADHD). *Educ Health* 2011;24:577.
- 20 Grewal P, Alagaratnam S. The quality and readability of colorectal cancer information on the Internet. *Int J Surg* 2013;11:410–3.
- 21 Vetter D, Ruhwinkel H, Raptis DA, et al. Quality assessment of information on bariatric surgery websites. *Obes Surg* 2018;28:1240–7.

- 22 McCool ME, Wahl J, Schlecht I, *et al.* Evaluating written patient information for eczema in German: comparing the reliability of two instruments, DISCERN and EQIP. *PLoS One* 2015;10:e0139895.
- 23 Vaona A, Marcon A, Rava M, *et al.* Quality evaluation of JAMA patient pages on diabetes using the ensuring quality information for patient (EQIP) tool. *Prim Care Diabetes* 2011;5:257–63.
- 24 Dormuth CR, Carney G, Taylor S, *et al.* A randomized trial assessing the impact of a personal printed feedback portrait on statin prescribing in primary care. *J Contin Educ Health Prof* 2012;32:153–62.
- 25 Castillo-Ortiz JD, Valdivia-Nuno J de J, Ramirez-Gomez A, 'et al'. Legibilidad, relevancia y calidad de la información en Español en la web para pacientes con artritis reumatoide. *Reumatol Clínica* 2017;13:139–44.
- 26 Bea-Muñoz M, Medina-Sánchez M, Flórez-García MT. Readability and internet accessibility of informative documents for spinal cord injury patients in Spanish. *An Sist Sanit Navar* 2015;38:255–62.
- 27 Jo JH, Kim EJ, Kim JR, *et al.* Quality and readability of Internet-based information on halitosis. *Oral Surg Oral Med Oral Pathol Oral Radiol* 2018;125:215–22.
- 28 Swain D. Producing patient information – how to research, develop and produce effective information resources. *Health Expect* 2003;6:274–5.
- 29 Raynor DK, Knapp P, Silcock J, *et al.* "User-testing" as a method for testing the fitness-for-purpose of written medicine information. *Patient Educ Couns* 2011;83:404–10.
- 30 King SR, McCaffrey DJ, Bouldin AS. Health literacy in the pharmacy setting: defining pharmacotherapy literacy. *Pharm Pract* 2011;9:213–20.