in 2.6%. At least 41% of the patients had been treated with PPIs for more than a year; for the rest of the patients, no prescription data were available prior to joining the pharmacy service. The mean number of drugs prescribed per patient was 8.3 ± 5 , 1.2% were being treated with an NSAID, 42.3% were taking acetylsalicylic acid concomitantly at antiplatelet doses and 3.8% were given bisphosphonates. The prevalence of patients being treated with a PPI and with a history of fracture was 48.7%.

Conclusion and relevance There was a high prevalence of patients without a clear indication for the prescription of PPIs in the GHC. This makes it necessary to review the treatments to assess possible deprescription of these drugs. In addition, their administration could be related to an increased risk of fractures due to its high prevalence.

REFERENCES AND/OR ACKNOWLEDGEMENTS

Conflict of interest No conflict of interest

5PSQ-121 FIDAXOMICIN RELATED METABOLIC ACIDOSIS: A CASE REPORT

¹B Sema Serrano, ²A Serrano Martínez, ¹A Valladolid Walsh*, ¹I Pérez Alpuente, ¹V Lerna Gaude, ¹M Clemente Andújar, ¹R Aldaz Francés. ¹Complejo Hospitalario Universitario De Albacete, Pharmacy, Albacete, Spain; ²Complejo Hospitalario Universitario De Albacete, Haematology, Albacete, Spain

10.1136/ejhp-pharm-2021-eahpconf.240

Background and importance Fidaxomicin is a macrolide antibiotic used to treat intestinal *Clostridium difficile* (CD) infection in the absence of metronidazole or vancomycin treatments. Pharmacovigilance collects information, and analyses and notifies cases of suspected adverse drug reactions (ADRs) to prevent them occurring in the future.

Aim and objectives To describe a case of metabolic acidosis in a patient treated with fidaxomicin and establish its possible association.

Material and methods We describe the case of an 82-year-old man diagnosed with multiple myeloma and treated with two full cycles of bortezomib-dexamethasone. He was referred to the emergency department after presenting with melanic diarrhoea for 1 week. As a result, he was hospitalised and diagnosed with upper gastrointestinal bleeding, acute prerenal renal failure, mild thrombopenia, hypokalaemia and hyponatraemia. After fluid and electrolyte stabilisation, it was decided to start fidaxomicin 200 mg/12 hours due to fever, confusional syndrome, persistence of diarrhoea and positive CD toxin test. The following constants were measured to confirm metabolic acidosis: gas level of bicarbonate (HCO_3^-), partial pressure of carbon dioxide (pCO_2), hydrogen ion potential (pH) and anion GAP. The degree of drug adverse reaction causality was evaluated using the Naranjo algorithm.

Results Two blood gas tests on consecutive days confirmed very low HCO_3^- (9 mmol/L) and pCO_2 (16 mm Hg) with normal pH (7.4), after which the patient was diagnosed with compensated metabolic acidosis with normal GAP anion. Finally, it was decided to suspend fidaxomicin and in the following days the patient experienced a progressive clinical improvement. Naranjo's algorithm established the causality relationship as 'probable' (score of 6). The regional pharmacovigilance centre (RPC) was notified.

Conclusion and relevance The European Medicines Agency's technical sheet for fidaxomicin does not describe metabolic acidosis as an ADR. However, UpToDate Clinical Library reports <2% of cases of metabolic acidosis in adults treated with fidaxomicin. The RPC reported this case as the only fidaxomicin ADR notified in our country.

REFERENCES AND/OR ACKNOWLEDGEMENTS

Conflict of interest No conflict of interest

5PSQ-122 HEALTH RESULTS AFTER INFlixIMAB PHARMACOKINETIC MONITORING IN INFLAMMATORY BOWEL DISEASE

¹M Herrera Expósito*, ¹JE Martínez De La Plata, ²D Rubio Calvo, ³CM Pinto Nieto, ¹J Canto Mangana, ¹MA Castro Vida. ¹Hospital De Poniente, Pharmacy Department, El Ejido-Almería, Spain; ²Hospital De Poniente, Hospital Pharmacy, El Ejido-Almería, Spain; ³Hospital De Guadix, Pharmacy Department, Guadix-Granada, Spain

10.1136/ejhp-pharm-2021-eahpconf.241

Background and importance Infliximab (IFX) is a monoclonal antibody (inhibitor of alpha tumour necrosis factor (anti-TNF α)) which is used in the management of inflammatory bowel disease (IBD). However, some patients do not show clinical benefit for this therapy or they show loss of response over time. For this reason, it is advisable to individualise and optimise the therapy through therapeutic drug monitoring (TDM).

Aim and objectives To analyse the clinical situation of patients according to proactive monitoring of serum levels of IFX in IBD and pharmacokinetic recommendations in their management.

Material and methods A prospective study was carried out in a 350 bed general hospital. Patients with IBD and IFX determinations with pharmaceutical interventions performed between January 2019 and July 2020 were selected. Data collected were: sex, median age (range), analytical parameters before intervention (mean (SD) albumin, C reactive protein (CRP), α 1-acid glycoprotein (AGP), faecal calprotectin (FC)) and clinical status before intervention (good general condition (GGC), regular general condition (RGC) and bad general condition (BGC)). Mean (SD) IFX levels and anti-IFX antibodies (ATI) were measured and patients with concomitant immunomodulatory treatment were registered. According to population studies, new interventions were recommended by the pharmacy service (intensification, swap, deintensification, non-valuable). Analytical parameters (CRP, AGP, FC) and clinical status of the patient 3 months after the intervention were analysed. Data were obtained from the assisted electronic prescription programme and digital medical records.

Results 55 patients with interventions were monitored, 38 men (69.1%), mean age 39 (20–70) years. Analytical parameters before the intervention were: albumin 4.04 (0.3) g/dL, CRP 0.78 (0.75) mg/dL, AGP 87.35 (29) mg/dL and FC 190.25 (148.61) $\mu\text{g/g}$. Clinical status before the intervention was GGC 9 (19.36%), RGC 30 (54.55%) and BGC 16 (29.11%). IFX levels were 3.04 (4.22) $\mu\text{g/mL}$ and ATI 1.48 (2.61) $\mu\text{g/mL}$. Patients with concomitant immunomodulatory treatment 32 (58.18%). All recommended interventions were accepted: intensification 41 (74.55%), swap 10 (18.18%), deintensification 3 (5.45%) and non-evaluable 1 (1.82%). Analytical parameters 3 months after the intervention were: CRP